

Comment on “Chiral Dilepton Model and The Flavor Question”

Recently in Ref. [1] a model was considered based on the

$$SU(3)_c \otimes SU(3)_L \otimes U(1)_X$$

gauge symmetry. This model is identical to a model previously proposed in Ref. [2, 3] and has precisely the same gauge group and representation content. Here we just list some features associated with this model and address the reader to Refs. [2, 3, 4] for details. In particular we correct some incorrect results and misconceptions of Ref. [1]. The model has extra gauge bosons with single and double charge Y^-, Y^{--} , called dileptons in Ref. [1] (V^-, U^{--} in Refs. [2, 3]) and also a new neutral gauge boson Z^0 . Both kinds of vector bosons gain mass mainly from the vacuum expectation value (VEV), w of the Higgs triplet ($\mathbf{3}, +1$) which precipitates the first symmetry breaking. Ref. [1], using LEP measurement of the quantity $1 - M_W^2/M_Z^2$, obtained the lower limit on the new gauge bosons:

$$M_{Z'} \geq 300 \text{ GeV} \quad M_Y \geq 230 \text{ GeV}. \quad (1)$$

The Z^0 boson couples to flavor changing neutral currents which give a contribution to the $K^0 - \bar{K}^0$ mass difference and this imposes the stronger limit on its mass given in Ref. [3]:

$$M_{Z'} > 40 \text{ TeV}. \quad (2)$$

This implies that w is of the order of 8 TeV and consequently

$$M_{Y^-, Y^{--}} > 4 \text{ TeV}. \quad (3)$$

On the other hand, taking into account the three triplets as well as the sextet needed in the model in order to give mass to all the fermions, we found that [4]

$$\frac{M_Z^2}{M_W^2} = \frac{1 + 4t^2}{1 + 3t^2}, \quad (4)$$

where $t = g_X^2/g^2$. However, in order to be consistent with experimental data Eq.(4) must be numerically equal to $1/\cos^2\theta_W$, where θ_W is the weak mixing angle in the standard electroweak model. This implies

$$t^2 = \frac{\sin^2\theta_W}{1 - 4\sin^2\theta_W}. \quad (5)$$

Ref [1] assumed a normalization for the $U(1)_X$ and T_3 generators, prompted by a potential unification theory, which imposes the condition

$$g_X^2/g^2 = 4/5. \quad (6)$$

However, from Eq. (5), and using $\sin^2\theta_W \simeq 0.23$ we obtain $g_X^2/g^2 \simeq 2.87$ which is in conflict with (6). Then, only the lower bounds, Eq. (2 and (3) on the masses of the new gauge bosons arise.

The parameter $\xi = (M_Z^2/M_{Z'}^2)_0$, where the subscript denotes “lowest order”, was used in Ref. [1] in order to study the corrections to the standard model predictions. It was found that

$$\xi = \frac{g^2(3 + 8t^2)}{4(1 + 2t^2)} \frac{v^2}{w^2}, \quad (7)$$

where v and U are, respectively, the vacuum expectation value of the triplet with $X = 0$ and $X = +1$. Since $w > 8 \text{ TeV}$ and as we must consider v is of the order of the Fermi scale we see that the ξ -parameter is too small to be measured in currently energies and the new bosons will be detected only indirectly at new generations high-energy colliders.

I thank O.F. Hernández and R. Foot for useful discussions. This work was partially supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

PACS numbers: 12.15.Cc; 14.80.-j

V. Pleitez

Instituto de Física Teórica

Universidade Estadual Paulista, Rua Pamplona, 145

01405-900–São Paulo, SP, Brazil.

References

- [1] P.H. Frampton, Phys. Rev. Lett., **69**, 2889(1992).
- [2] F. Pisano and V. Pleitez, “Neutrinoless double beta decay and doubly charged gauge bosons”, IFT-P.017/91, July 1991.
- [3] F. Pisano and V. Pleitez, Phys. Rev. D**46**, 410(1992).
- [4] R. Foot, O. F. Hernandez, F. Pisano and V. Pleitez, Leptons masses in an $SU(3)_L \otimes U(1)_N$ Gauge Model, preprint IFT-P.025/92, McGill/92-31, submitted for publication.